

**CS/B.TECH(EE/EEE/EIE/ICE-New)/SEM-4**

**/PH(EE)-401/2012**

**2012**

**PHYSICS-II**

Time Allotted : 3 Hours

Full Marks : 70

*The figures in the margin indicate full marks.*

*Candidates are required to give their answers in their own words  
as far as practicable.*

**GROUP – A**

**(Multiple Choice Type Question)**

1. Choose the correct alternatives for any *ten* of the following:

10 x 1 = 10

- i)  $\text{He}^3$  and muon are
  - a) fermions
  - b) bosons
  - c) fermions and bosons respectively
  - d) bosons & fermions respectively.
- ii) The degrees of freedom for a system of N particles with K constraint relations is given by
  - a)  $N-K$
  - b)  $N-3K$
  - c)  $3N-K$
  - d)  $3K-N$ .
- iii) The coordination number for FCC structure is
  - a) 6
  - b) 8
  - c) 12
  - d) 5
- iv) The dielectric constant for a conductor is
  - a) 0
  - b) 1
  - c) -1
  - d) infinity

- v) Fermi-Dirac distribution approaches Maxwell-Boltzmann distribution at
- low temperature & high density
  - high temperature & low density
  - low temperature & low density
  - high temperature & high density
- vi) If  $E_1$  is the energy of the ground state of a one-dimensional potential box of length  $l$  and  $E_2$  be the energy of the ground state when the length of the box is halved, then
- $E_2 = 2E_1$
  - $E_2 = E_1$
  - $E_2 = 4E_1$
  - $E_2 = 3E_1$
- vii) The reciprocal lattice of a body centered cubic (*bcc*) lattice is
- bcc*
  - fcc*
  - sc*
  - hcp*
- viii) The wave function of a particle is  $\psi = A \cos^2 x$  for  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ . Then, the value of A is
- $\sqrt{\frac{8}{3\pi}}$
  - $\sqrt{\frac{3}{8\pi}}$
  - $\sqrt{\frac{1}{2\pi}}$
  - $\sqrt{\frac{3}{2\pi}}$
- ix) The density of free electron states in a metal varies with energy E as
- $\sqrt{E}$
  - $E^2$
  - $E^0$
  - $\frac{1}{E}$

- x) Curie-Weiss law is obeyed by
- paramagnetic materials
  - anti-ferromagnetic materials
  - ferromagnetic material above the Curie temperature
  - ferromagnetic materials below the Curie temperature.
- xi) The Miller indices of a plane parallel to XY plane is
- (100)
  - (010)
  - (001)
  - (110)
- xii) If  $\sigma$  and  $k$  be the electrical and thermal conductivities in a solid, then according to Wiedemann-Franz law.
- $\frac{\sigma}{kT} = \text{const.}$
  - $\frac{k\sigma}{T} = \text{const.}$
  - $\frac{k}{\sigma T} = \text{const.}$
  - $\sigma kT = \text{const.}$
- ( Where T is the temperature )
- xiii) The product generalized ( $Q_1$ ) and generalized displacement ( $\delta q_j$ ) must have the dimension of
- force
  - work
  - power
  - length.
- xiv) The spacing between the  $n$ th energy state and next energy state in a one-dimensional potential box increase by
- $2n-1$
  - $2n+1$
  - $n-1$
  - $n+1$
- xv) In an n-type semiconductor, donor level
- is nearer to conduction band
  - is at the middle between valence and conduction
  - is nearer to valence band
  - is not formed at all.

## GROUP – B

### (Short Answer Type Questions)

Answer any *three* of the following.

3 x 5 = 15

2.
  - a) Describe briefly micro-state and macro-state with suitable examples.
  - b) Show that the average energy of electrons at  $T=0$  K is given by  $\frac{3}{5} E_F$  ( where  $E_F$  is the Fermi energy ). 2+3
3.
  - a) What do you mean by cyclic coordination? Explain with an example.
  - b) Show that if a given coordination is cyclic in Lagrangian, it will also be cyclic in Hamiltonian. 2+3
4.
  - a) Define atomic polarizability. Establish a relation between polarization and atomic polarizability.
  - b) Calculate the induced dipole moment per unit volume of the He gas if it is placed in an electric field of  $6000 \text{ V cm}^{-1}$ . The atomic polarizability of He is  $0.18 \times 10^{-40} \text{ Fm}^2$  and density of He is  $2.6 \times 10^{25} \text{ atoms per m}^3$ . 3+2
5.
  - a) Derive Curie's law of paramagnetism in the framework of Langevin's theory.
  - b) Are all orientations of the magnetic dipoles possible in quantum theory? Explain. 4+1
6.
  - a) Explain what you mean by degeneracy of an eigenstate with an example.
  - b) The eigenvalue equation for the momentum operator is 
$$\left(\frac{\hbar}{t}\right) \left(\frac{\partial \psi}{\partial x}\right) = \lambda \psi.$$

Solve the above equation and hence show that for  $\psi$  to be a physically admissible eigenstate, the eigenvalue  $\lambda$  must be real. 2+3
7. Derive the Bragg's law of X-ray diffraction from Laue equation and deduce the vector form of Bragg's law of X-ray diffraction in reciprocal space. 2+3

## GROUP – C

### (Long Answer Type Questions)

Answer any *three* of the following.

3 x 15 = 45

8. a) A free particle of mass  $m$  is confined within  $x = 0$  and  $x = L$ .
- (i) Write down Schödinger time-dependent equation for such a system.
  - (ii) Solve the equation to find out the normalized eigenfunctions.
  - (iii) Show that the eigenfunctions corresponding to two different eigenvalues are orthogonal. 1+4+3
- b) If  $\hat{P}$  and  $\hat{L}$  be the momentum and angular operators, find the values  $[\hat{L}_x, \hat{x}]$  and  $[\hat{L}_x, \hat{y}]$ . 2+2
- c) Find the expectation value of  $x$  for the wave function given by  $\psi(x) = Ae^{-bx^2}$ . 3
9. a) The energy wave vector dispersion relation or a one-dimensional crystal of lattice constant  $a$  is given by  $E(k) = E_0 - \alpha - 2\beta \cos(ka)$ , where  $E_0, \alpha, \beta$  are constants.
- i) Find the value of  $k$  at which the value of an electron is maximum.
  - ii) Find the difference between the top and the bottom of the energy band.
  - iii) Obtain the effective mass  $m^*$  of the electron at the bottom and at the top of the band. 2+2+2
- b) What do you mean by density of states? Show that the density of states of free electrons vary with energy  $E$  as  $\sqrt{E}$ . 1+4
- c) In sodium metals, the free electron density is  $2.5 \times 10^{28} \text{ m}^{-3}$ . Calculate the Fermi energy and the fermi temperature. 2+2
10. a) Define Hamiltonian of a dynamical system. When does it represent the total energy of the system? Explain. 2+3

- b) The Lagrangian of a particle of mass  $m$  in one dimension is given by

$$L = \frac{1}{2} m(\dot{x}^2 - \omega^2 x^2)e^{bt}$$

Obtain the canonical momentum and equation of motion. Is the Hamiltonian constant of motion? 3+3

- c) Deduce D'Alembert's principle from the principle of virtual work. 4
11. a) What do you mean by symmetric and anti-symmetric wave function? How does Fermi-Dirac(FD) statistics differ from Bose-Einstein(BE) statistics? 2
- b) Explain graphically the Fermi distribution at zero and non-zero temperature. 3
- c) Derive Planck's radiation law from BE statistics. State clearly the assumptions made in the theory. 3+2
- d) Compute the specific heat of a free electron gas using classical statistics. Using FD statistics, argue that the specific heat of electrons should vary linearly with temperature(T). 2+3
11. a) What is Larmor frequency? 2
- b) With the help of Weiss molecular field theory of ferromagnetism, derive the Curie-Weiss law. 5
- c) Draw the B-H curve for a ferromagnetic material and identify the retentivity and the coercive field on the curve. What is the energy loss per cycle? 3+1
- d) Explain the reason behind the negative susceptibility of diamagnetic material. 2
- e) Calculate the Bohr magneton for  $Gd^{+3}$ . The electronic configuration for  $Gd^{+3}$  is  $4f^2 5s^2 5p^6$ . 2

=====